



Highlander Help

This site was created for *you*

Here you will find answers, explanations, and resources to guide you through the homework and develop a better understanding of the material

You deserve to succeed and to have more time to do the things you love to do.

Find this material useful?

*You can help our team to keep this site up and bring you even more content—**consider donating** via the link on our site.*


Still having trouble understanding the material?

*Check out our “**Tutoring**” page to find the help you need.*

Good Luck!

1. Two masses $M_1 = 2 \text{ kg}$ and $M_2 = 4 \text{ kg}$ are attached by a string as shown. They start from rest and move with no friction until they reach a velocity of 6.5 m/s. When do they reach that speed, in s?

- a. 0.68
- b. 4.3
- c. 2.0
- d. 1.4
- e. 3.3



$\cancel{v} + \frac{1}{2}at^2$

$(2) = at$
207

$g = (2+4)a$

$3.267 = g = ma$

$\frac{2(9.8)}{6} = a$

$2(9.8)$
 mg

$W = (F) \cdot D$

$W = \frac{1}{2}mv^2 - \frac{1}{2}mv_0^2$

$W = \frac{1}{2}(2)(6.5)^2$

$W = 42.25$

$2(g)D = \frac{1}{2}(2)(6.5)^2$

$a = 3.267$


$2077 = \frac{1}{2}at^2$

$= \frac{1}{2}(\frac{2 \cdot 9.8}{6})t^2$

$X = \frac{1}{2}(\frac{9.8}{3})t^2$

2. "A light fixture of mass 3.55 kg hangs by two wires (arranged like a "Y"), each of which makes an angle of 10 degrees with the ceiling. What is the tension, in N, in one of the wires?"

- a. 200
- b. 100
- c. 25
- d. 20
- e. 10



$T_1 \cos 10 - T_2 \cos 10 = 0$

$T_1 \sin 10 + T_2 \sin 10 = 3.55$


$2 \times \sin 10 = 3.55$

$\frac{100}{1.775} = \frac{3.5}{x}$

3. A light fixture is suspended from a wall and a ceiling by wires, as shown. The tension, T_1 in wire 1 is 3.5 N. What is the mass, M , in kg?

- ~~a. 3.4~~
- ~~b. 2.0~~
- ~~c. 1.7~~
- ~~d. 0.64~~
- e. 0.21

$T_1 = 4$



$3.5 \cos 90 - T_2 \cos 30 = 0$


$T_2 \sin 30 + 3.5 \sin 90 = 9.8x$

$T_2 \cos 30 = 3.5 \cos 20$

$T_1 \sin 30 + 3.5 \sin 90 = M(9.8)$

4. Two blocks are on a horizontal, frictionless table. A force of 2.0 N pulls a block that has $m=3.0\text{ kg}$. The block is connected to a second block, $M=4.0\text{ kg}$, by a wire. What is the tension, in N , in the wire?

- a. 1.1
- b. 0.86
- c. 0.29
- d. 2.0
- e. 1.0



$$2 = 7a$$

$$a = \frac{2}{7}$$

$$F = m \left(\frac{2}{7} \right)$$


$$\frac{8}{7}$$

$$2 = \dots \quad \frac{2}{4} (2)$$

5. A beam is tilted up by 0.02 m at one end, and it is 1 m long. Friction is insignificant. Mass of the glider is 1 kg . What is the magnitude of the component of net force, in N , acting on the glider along the beam.

- a. 100
- b. 2
- c. 5
- d. 0.02
- e. 0.2

soh / cah / toa



$$mg \sin(1.145942) \quad | \quad \sin \theta = \frac{.02}{1}$$

1/96

6. An elevator of mass 1000 kg pulls down on one side of a cable that goes over a pulley that has no friction. A counter weight of 900 kg pulls on the other side. The elevator starts to fall with no friction. What is the net force on the system, in N ? Take the elevator's direction of motion as positive.

- a. 50
- b. 980
- c. 100
- d. 460
- e. 20


$$1000 - 900$$

$$100 (9.8)$$

$$mg = (1000 - 900)g (9.8)$$

7. On a Force table, forces are applied to a small ring near the center. If the forces are $F_1 = -0.9i + 0j$ and $F_2 = 0i - 0.75j$, what is the magnitude, in N, of a third force which will keep the ring in equilibrium without touching the pin at the center?


- a. 1.2
- b. 1.6
- c. 1.4
- d. 0.9
- e. 0.6



8. A crate of weight 100-N is sitting on a ramp with a 30 degree slope, as shown. The playful man lets go and the crate slides down with no friction and an acceleration a_1 . He then places another crate of half the weight on the ramp. Again, he lets go and the second crate slides down with acceleration a_2 . What are values of a_1 and a_2 in m/s^2 ?

- a. 9.8, 4.9
- b. 4.9, 2.5
- c. 9.8, 9.8
- d. 4.9, 9.8
- e. 4.9, 4.9

Handwritten equations for problem 8:
 $m = a$
 $F = 100 \sin 30$
 $F = 50 \sin 30$
 4.9
 4.9



9. The man lets go of the rope attached to the 100-N crate. It starts from rest and slides down the 30-degree slope with no friction. The crate's velocity increases to 9.8 m/s at a time t . What is t in s?

- a. 9.8
- b. 0.01
- c. 0.3
- d. 2.0
- e. 9.8

Handwritten equations for problem 9:
 $\frac{1}{2} m v^2 = 100 \sin 30$
 $489.5 = 100 \sin 30 D$
 $9.8 = D$
 $9.8 = v + \frac{1}{2} a t^2$

Handwritten notes for problem 9:
 $100 \sin 30$
 $10,204$
 $F = m \cdot a =$
 $a = 4.9$

10. A CONSTANT force acts on an object and increases its kinetic energy, by 24 J. The object moves from $(7\mathbf{i} - 8\mathbf{j} + 4\mathbf{k})\text{m}$ to $(11\mathbf{i} - 5\mathbf{j} + 4\mathbf{k})\text{m}$. The net force acting on the object is equal to $(F_x\mathbf{i} + 4\mathbf{j} + 5\mathbf{k})\text{N}$. What is F_x , in N.

- ~~a. 1~~
- b. -4
- c. 4
- d. 3**
- e. -3

$$24 \text{ J} = (F, 4, 5) \cdot (4, 12, 3)$$

$$24 = 4F + 12 + 0$$

$$\begin{matrix} 11 & - & 7 & & 4 \\ - & 5 & + & + & 8 & & 3 \\ & & & & 4 & - & 4 & & 0 \end{matrix}$$

+ 4 (x)	4x	
+ 3 (y)	12	= 24
+ 0 (z)	0	3 = x

11. As shown in the figure, a block is pushed up against a vertical wall by a force 20 N. The force is at an angle of 40 degrees from horizontal. The coefficient of static friction between the block and the wall is 0.50. Find the maximum mass, in kg, that the force can prevent from sliding down.

- ~~a. Infinite~~
- b. 0.92
- c. 2.1**
- d. 0.52**
- e. 10

$$20 \cos 40 = F_N = 15.321$$

$$F_f = 15.321(0.5)$$


$$7.6605$$

$$20 \sin 40 = F_y = 12.856$$

$$F_A = 20.91619 = m(9.8)$$

$$F_N = 20 \cos 40$$

$$F_f = .5 (20 \cos 40)$$




mg

$$9.9m = 10 \cos 40 =$$

12. A block sits on a horizontal surface with a coefficient of static friction of 0.2 between them. A horizontal force of 14 N is just able to move the block parallel to the surface. What is the mass, in kg, of the block?

- a. 7.1
 b. 14
 c. 70
 d. 1.4
 e. 3.5




$$14\text{ N} - 0.2\text{ mg} = 0$$

$$14 = 1.96\text{ m}$$

13. Two masses are attached by a string as shown. The mass, M_2 , is 4 kg and the coefficient of friction is 0.5. What is the maximum mass M_1 , in kg, that allows the system to stay at rest?

- a. 0.3
 b. 2
 c. 4
 d. 0.5
 e. 8

$$F = F$$




$$\mu m g = m g$$

$$\mu 4 g = 2$$

$$0.5 4 g = 2$$


14. The block is at rest. Then, the ramp is gradually tilted. At an angle of 14 degrees, the block begins to slide. What is the coefficient of static friction between the block of unknown mass, m , and the ramp?

- a. Can't tell; need m
 b. 0.87
 c. 0.61
 d. 0.55
 e. 0.25



15. A woman pulls a block along a horizontal surface at a constant speed with a 15-N force acting 20° above the horizontal. She does 85 J of work. How many meters does the block move?

- a. 5.6
- ~~b. 90~~
- c. 6.0
- ~~d. 0.16~~
- e. 3



$$85 = 15 \cos 20 \cdot \text{D}$$

16. A person does 200 J work lifting an object from the bottom of a well at a constant speed of 2.0 m/s in a time of 5.1 s. What is the object's mass? (Neglect friction.)

- a. 20
- b. 2.0
- c. 6.2
- d. 2.1
- e. 4.0

$$200 \text{ J} = 10.2 \cdot m \cdot g$$

$\frac{2 \text{ m}}{5} \cdot 5.1$

17. A woman throws a 2.0-kg ball from the origin to a point at $(20 \text{ i} + 3 \text{ j} + h \text{ k})$ meters, where k is the upward unit vector. The work done by the gravitational force on the ball is -290 J. What is the height, h ?


- a. 19
- b. 15
- c. 39
- d. 7
- e. 150

$$-290 \text{ J} = 2 \cdot (9.8) \cdot h$$

Version B

8. Suddenly, the driver of a fast car travelling 50 m/s sees a deer and slams on the brakes. The car travels for 10 s before it stops. What is the coefficient of kinetic friction between the tires and the road?

- a. 0.13
- b. 0.26
- c. 0.32
- d. 0.41
- e. 0.51



$$\frac{50 \text{ m}}{10 \text{ s}} = F$$

$$F_f = mg\mu$$

$$50 \text{ m} - \frac{m}{m} \mu (10) = 0$$

$$0 = 50 - 10a$$

$$a = -5 \text{ m/s}^2$$

$$50 \text{ m}$$

$$+\frac{1}{2}(50)^2 = 500g\mu$$

$$W = -\frac{1}{2}(50)^2 \text{ m/s}^2 = M \text{ m/s}^2 \cdot 5 \text{ m/s}^2$$

$$F_g = mg$$

$$500 = \frac{1}{2}mv^2$$


$$\frac{50}{2} = mg$$

$$50 = 98\mu$$

$$\mu = .51$$

9. An object falls vertically downward in water at a constant speed. The viscosity of the water does work of -20 J as the object falls 0.80 m. What is the mass, in kg, of the object?

- a. 2.0
- b. 20
- c. 3.7
- d. 2.6
- e. 1.7



$$-20 = F \cdot D$$

$$-20 = mg \cdot 0.8$$

10. A constant force of $(2\mathbf{i} - 15\mathbf{j} + 2\mathbf{k})$ N acts on a particle as it moves from the origin to a point $(4\mathbf{i} + 3\mathbf{j} + 5\mathbf{k})$ m. How much work, in J, does the force do during this displacement?

- a. +30
- b. -27
- c. +45
- d. -45
- e. +37

$$W = F \cdot D$$

$$W = (2, -15, 2) \cdot (4, 3, 5)$$

$$8 - 45 + 10$$